Supplementary information, Fig. S6. In this figure, cancer cells were treated with DM-αKG (15 mM) for 6 hours to determine the ROS level, or 24 hours to assess pyroptotic features (including cell morphology, GSDMC cleavage, LDH release, and Annv⁺/PI⁺ cells), unless specially indicated otherwise.


- (a) DM-αKG-induced ROS level and pyroptosis were not influenced by dimethyloxalylglycine (DMOG). HeLa cells were pretreated with DMOG (50 μM) for 2 hours before DM-αKG treatment.
- (b) The knockdown efficiency of OGDH, IDH1, IDH2, MDH1, MDH2 and LDHA in HeLa cells, as determined by western blot or RT-qPCR analysis.
- (**c**, **d**) Knocking down IDH1 or IDH2 showed no effects on DM-αKG-induced pyroptotic morphology, LDH release (c), or ROS levels (d) in HeLa cells.
- (e, f, g, h) Knocking down OGDH increased ROS levels (e), GSDMC cleavage and LDH release (f), Annv⁺/PI⁺ cells (g) and pyroptotic morphology (h) as induced by DM-αKG. HeLa cells were treated with DM-αKG with or without OGDH knocked down.
- (i) Knocking down LDHA had no effect on DM-αKG-induced ROS levels or pyroptotic morphology in HeLa cells.
- (j) Knocking down MDH2 had no effect on DM-αKG-induced ROS levels or pyroptosis in HeLa cells, as determined by cell morphology and LDH release.


- (k) Knocking down MDH1 impaired DM- α KG-induced pyroptosis in SGC-7901 (top) and B16 (bottom) cells, as determined by cell morphology, GSDMC cleavage and LDH release.


 (l) The expression levels of MDH1^{WT} and MDH1^{H187Y} in HeLa cells.
- (m, n, o) Knocking down caspase-8 (m), DR6 (n) or GSDMC (o) impaired Octyl-L-2HG-induced pyroptosis in HeLa cells, as determined by cell morphology and LDH release.
- (p, q) Knocking out DR6 (p) and GSDMC (q) abolished hypoxia-induced pyroptosis in HeLa cells.


Tubulin was used to determine the amount of loading proteins. All data are presented as the mean \pm SEM of two or three independent experiments. *** p<0.001, ns: not significant. The data were analyzed using two-way ANOVA followed by the Bonferroni test.

Supplementary information, Figure S6









g

